

Analysis of Intermediate-Energy (γ, n) Reactions in Complex Nuclei*

H. G. DE CARVALHO, M. FOSHINA and J. D. PINHEIRO FILHO**

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro RJ

and

V. DI NAPOLI, J. B. MARTINS***, F. SALVETTI, O. A. P. TAVARES***and
M. L. TERRANOVA

Istituto di Chimica Generale ed Inorganica dell'Universita, Roma

Recebido em 12 de Agosto de 1977

An analysis of the available experimental data on (γ, n) reaction cross sections, at intermediate energies, was made in order to obtain their dependence on the mass number A_t . The results were compared with those obtained by means of a Monte Carlo calculation based on photon-initiated intranuclear cascades. A relation has been established between the cross sections of elementary processes (pion photoproduction on single nucleon and Levinger's quasi-deuteron photodisintegration) and the (γ, n) probability. Cross section formulas have been obtained which give an A_t -dependence of the type aA_t^b , with $b = 0.86$ and a approximately equal to the cross sections of the elementary processes $\gamma + n \rightarrow n + \pi^0$ and $\gamma + p \rightarrow n + \pi^+$. These formulas reproduce almost all known experimental cross sections within a factor of 2, which seems to be a rather good accuracy

*Based on part of the Doctoral Thesis presented by J.B.Martins at the *Centro Brasileiro de Pesquisas Físicas*, Rio de Janeiro, November 1974. This work has been supported in part by the Brazilian *Comissão Nacional de Energia Nuclear*, Brasil, and by the *Consiglio Nazionale delle Ricerche*, Italia.

Permanent address: *Instituto de Física, Universidade Federal Fluminense*, Niteroi, RJ.

Permanent address: *Centro Brasileiro de Pesquisas Físicas*, Rio de Janeiro, RJ.

in view of the large range covered by both cross sections values and target mass numbers.

Uma análise estatística das seções de choque experimentais de reações (γ, n) , em energias intermediárias, foi efetuada com a finalidade de estabelecer uma dependência com o número de massa, do núcleo alvo, A_t . Os resultados foram comparados com os obtidos por meio de um cálculo de Monte Carlo baseado em cascatas intranucleares iniciadas por fótons. Foi estabelecida uma relação entre as seções de choque dos processos elementares (fotoprodução de mésons a partir de nucleons isolados e a foto-desintegração do quase-deuteron) e a probabilidade de reações (γ, n) . Foram determinadas fórmulas que fornecem uma dependência das seções de choque com A_t , do tipo aA_t^b , sendo $b = 0,86$ e a aproximadamente igual às seções de choque dos processos elementares $\gamma + n \rightarrow n + \pi^0$ e $\gamma + p \rightarrow n + \pi^+$. Essas fórmulas reproduzem bem as seções de choque experimentais dentro de um fator 2, o que pode ser considerado um bom resultado, em vista dos grandes intervalos varridos pelas seções de choque e pelo número de massa do núcleo alvo.

1. INTRODUCTION

Above about 0.1 GeV, photons interact with nuclei giving rise to different kinds of reactions such as direct reactions, fragmentation, spallation and fission.'

Direct reactions originate from the ejection from the struck nucleus of a very small number of nucleons and/or pions leading to a residual nucleus with an excitation energy not sufficient to "evaporate" other particles. Such reactions may be ascribed to a fast process that does not affect the rest of the nucleus. The (γ, n) and (γ, p) reactions are among the most remarkable examples of these processes.

In the region of intermediate energies (0.1-1.0 GeV), two models have successfully been proposed in order to explain the photon-nucleus interaction pattern: Lvinger's quasi-deuteron¹ and the photomesonic² interaction mechanisms. In the quasi-deuteron model, the photon is absorbed by a neu-

tron-proton pair within the nucleus and the cross section of most processes is strictly related to that of the deuteron photodisintegration, σ_d , and to the number of "quasideuterons" in the target nucleus. The quasideuteron contribution is important for energies up to 0.5 GeV, and becomes negligible above this energy value, due to the smallness of the cross section of the deuteron photodisintegration.

At incident energies higher than or equal to the threshold of π -meson photoproduction on single nucleons (about 0.15 GeV), the experiment shows that the slopes of the yield curves become steeper the higher the energy of the incoming photon is, and broad resonances have been observed corresponding to the resonances found for single and double-pion photoproduction. Another mechanism of absorption has to be invoked, therefore, which is related to the pion photoproduction. Photonuclear cross sections must hence be connected with the total inelastic cross section, σ_N , of the γ -nucleon interaction.

The photomesonic model suggests, as the primary interaction, the photoproduction of a real or virtual π -meson from a single nucleon, followed by two-step cascade-evaporation processes²⁻⁵. This mechanism becomes dominant as the energy increases.

Direct reactions, such as (γ, n) , represent a rather conspicuous contribution to the total γ -nucleus inelastic cross section. As a consequence, knowledge of the dependence of the (γ, n) cross section on such parameters as photon energy and target mass number can give valuable information about some scarcely known aspects of photon absorption, and clarify, to some extent, the mechanism of the interaction of photons with complex nuclei.

Since the experiment made by Baldwin and Klaiber⁶ in 1948 using *Bremstrahlung* from the General Electric 100-MeV Betatron to measure the yields of $^{12}\text{C}(\gamma, n)$ \rightarrow C and $^{63}\text{Cu}(\gamma, n)$ \rightarrow ^{62}Cu reactions. many attempts have been made to measure photoneutron cross sections covering wide ranges of either energies or mass number of target nuclei.

The present status of measurements of single-neutron photoproduction cross section is characterized by a striking difference in the number of expe-

periments, those made at lower energies being considerably more numerous than at intermediate and higher energies. Moreover, the use of almost monochromatic radiation produced by annihilation in flight of positrons, and of monochromatic gamma-rays from particle-induced reactions, has provided an important contribution to the understanding of the low-energy processes, since it allows for cleaner experiments than does the use of *Bremsstrahlung*. Excellent review articles have collected all known data on photoneutron cross sections in the giant resonance region⁷⁻⁹.

Unfortunately, the same is not true of the high-energy reactions, for which almost-monochromatic beams have not yet been extensively used for a number of reasons.

From all these facts, it is readily understood how relatively poor is the information presently available on (γ, n) reactions, at energies above 0.1 GeV, for both the scarcity of experimental data and the considerable discrepancy which exists between the results from different laboratories. As far as this conflict is concerned, one should keep in mind that the cross sections given by the different authors are in general mean cross sections calculated from *Bremsstrahlung* yield curves over wide energy ranges; and the difficulties inherent in using a *Bremsstrahlung* spectrum to excite a nuclear reaction very often prevent one from observing a finer structure of these curves. On the other hand, whatever the method of computation, small changes in the yield curve can introduce large fluctuations in the calculated cross sections¹⁰.

A number of papers have been published¹¹⁻³³ on (γ, n) reactions on light and more complex nuclei in the energy range 0.1-1.0 GeV. In spite of the above-mentioned discrepancies, a general trend of increasing cross sections with increasing mass number of the target nucleus can be deduced from the analysis of the data published in the literature from 1964 onward.

Some effort has been made to relate the (γ, n) cross section to that of pion photoproduction. De Carvalho and coworkers¹⁶ attempted to find a connection between the number of neutrons of the target nucleus, the cross section of pion photoproduction, and the transparencies of the nucleus to pions and recoil nucleons. Calculated values of nuclear transparen-

cies^{34,35}, however, did not entirely corroborate this point of view, especially in the case of medium-weight and heavy nuclei, and the conclusions was drawn that the cross sections calculated in this way represent only lower limits for the (γ ,n) reactions.

Researchers at Lund^{36,37} were able to relate the (γ ,n) cross section to the number of loosely bound neutrons in a few upper shell ("valence neutrons") in the nucleus. However, their calculations were restricted to a small number of light nuclei only.

In recent years, Monte Carlo calculations have been used in the study of (γ ,n) reactions. One of these calculations covers the *Bremsstrahlung* energy range 40-350 MeV, Ref.38, making it thus impossible a comparison with data from experiments at higher energies; the other³⁹ is restricted to one target nucleus only (¹⁹⁷Au) as far as the (γ ,n) reaction is concerned, and the calculated yield does not seem to reproduce satisfactorily enough the experimental results.

The decision as to which method of adjustment and/or comparison of the experimental data is preferable depends on whether the method chosen is suitable in providing the most general understanding of the phenomenon under investigation or not. Also, it should be able to give rather detailed information about the mechanism effective in producing the reactions considered. Furthermore, a test of its suitability might be the accuracy with which a cross section of a (γ ,n) reaction for any element can be calculated.

We believe that a very useful method would consist in handling the available data on (γ ,n) reaction in a statistical way and then, from the gross trend thus deduced, inferring some relationships between nuclear parameters and γ -nucleus interaction cross section on the basis of theoretical models.

Also, the aim of this work was to carry out a Monte Carlo calculation on the cross sections of (γ ,n) processes for target mass numbers between 12 and 238, at photon energies from 0.2 GeV up to 1 GeV, and to compare the experimental cross sections with the calculated ones.

It is to be hoped that a study along these lines will contribute to our understanding of direct reactions.

2. STATISTICAL TREATMENT OF THE AVAILABLE DATA

A total of 29 noteworthy cross section measurements are available to date for the (γ, n) reaction at energies between 0.1 GeV and 1.0 GeV. They take 14 different target nuclei, rather well distributed throughout the periodic table.

In some cases, only *Bremsstrahlung* activation yield curves have been published; therefore, the absolute cross sections (per photon) have been calculated by drawing least squares fits through the experimental yield values and by assuming a pure $1/k$ dependence of the *Bremsstrahlung* spectra upon the photon energy k .

Before entering into a more detailed discussion on the experimental data and their arrangement, we believe it worthwhile to make some specifications about the notation we have used. Yields are expressed as cross sections per equivalent quantum and denoted by the symbol σ_Q . The σ_Q values are functions of the *Bremsstrahlung* end-point energy E_0 . The cross section per photon, σ_k , also denoted as "absolute cross section", depends on the photon energy k .

The relation between $\sigma_Q(E_0)$ and $\sigma_k(k)$ is given by the integral equation

$$\sigma_Q(E_0) = \frac{1}{Q} \int_0^{E_0} \sigma_k(k) n(k, E_0) dk \quad , \quad (1)$$

where Q is the number of "equivalent quanta" (which represents the number of fictitious photons we should have if all the incident energy were equally distributed between photons of the maximum energy E_0), and $n(k, E_0)$ is the spectral density distribution of the *Bremsstrahlung* used. Since we are interested in the measurement of the mean values of σ_k in the energy range from about 0.1 GeV up to about 1.0 GeV (in most cases 0.3-

1.0 GeV), $n(k, E_0)$ can be approximated by Q/k and, consequently, the mean cross section $\bar{\sigma}_k$ in the energy range $E'_0 - E''_0$ can be expressed by

$$\bar{\sigma}_k = \frac{\sigma_Q(E''_0) - \sigma_Q(E'_0)}{\log(E''_0/E'_0)}, \quad (2)$$

$E'_0 - E''_0$ being an energy interval within which the trend of σ_Q versus $\log E_0$ may be approximated by a straight line.

In evaluating the mean cross sections per photon, $\bar{\sigma}_k$, in such a manner, one disregards, in some way, the background contribution to the measured yields due to the giant resonance region, which in the case of (γ, n) reactions is relevant. This contribution strongly depends on the actual shape of the Bremsstrahlung used, and thus different experimental arrangements may lead to remarkable different values of the measured σ_Q cross sections. Besides, the exact knowledge of the shape of the (γ, n) cross section resonance curves in the low energy region is not so straightforward, since in most cases the reported curves (see, for instance, Ref. 9) are the sum of (γ, n) and (γ, pn) processes. Also, the quasideuteron contribution from the end of the giant resonance up to about 70 MeV, is hardly evaluable. As a consequence, the low-energy tail contributes to an extent which is very difficult to be evaluated. We decided, therefore, to disregard such corrections and use Eq. (2) in calculating the mean cross sections in the energy range 0.3-1.0 GeV. Of course, the cross sections thus obtained are slightly overestimated, as will be proved by the results of the Monte Carlo calculation.

Table 1 reports the experimentally determined mean cross sections for each target nucleus, with the energy range covered by the measurements.

As a first approach, all these data were plotted against the target mass number, A_t , on a log-log graph and a straight line fitted to them by means of the least squares method¹⁰. Only two points were disregarded in this fit, in view of their very large deviation from the mean, namely, the cross sections relative to ^{14}N and ^{103}Rh . The extremely small value of the cross section of the $^{14}\text{N}(\gamma, n)^{13}\text{N}$ reaction finds an explanation in the low proton separation energy of ^{13}N (about 1.9 MeV) which makes ^{13}N very

Table 1 - Mean Experimental Cross Sections for (γ,n) Reactions at Intermediate Energies

Target Nucleus	Energy Range (GeV)	Laboratory	Cross Section (mb)	Ref.
^{12}C	0.2 - 0.72	INS, Tokyo	0.85 ± 0.10	12
	0.3 - 1.4	PTI, UAS	0.4 ± 0.1	13
	0.3 - 1.0	Frascati	1.18 ± 0.20	15
	0.2 - 1.0	Lund	0.6 ± 0.1	21,27
	0.3 - 1.0	Frascati	0.9 ± 0.1	32
^{14}N	0.32- 0.62	Lund	0.11 ± 0.05	23
^{16}O	0.3 - 0.8	Lund	0.6 ± 0.1	23
	0.3 - 1.0	Frascati	1.27 ± 0.20	29
^{19}F	0.3 - 0.9	Lund	0.42 ± 0.10	23
	0.3 - 1.0	Frascati	1.33 ± 0.20	24
	0.3 - 1.0	Frascati	1.30 ± 0.10	30
^{23}Na	0.3 - 1.0	Frascati	1.60 ± 0.20	30
^{31}P	0.3 - 0.8	Lund	0.9 ± 0.1	23
	0.3 - 1.0	Frascati	1.6 ± 0.1	33
^{52}Cr	0.3 - 1.0	Frascati	2.5 ± 0.5	33
^{55}Mn	0.1 - 0.8	Lund	7.7 ± 0.5	17
	0.3 - 1.0	Frascati	4.2 ± 0.6	31
	0.3 - 1.0	Frascati	3.5 ± 0.5	33
^{59}Co	0.2 - 0.8	Lund	3.3 ± 0.5	18
	0.3 - 1.0	Frascati	3.0 ± 0.5	33
^{75}As	0.3 - 0.9	Lund	8.7 ± 1.0	22
	0.3 - 1.0	Frascati	4 ± 1	33
^{103}Rh	0.4 - 0.9	Orsay	13 ± 5	19
^{127}I	0.15 - 0.80	Lund	10 ± 6	14
	0.1 - 0.8	Lund	4 ± 2	20
	0.3 - 1.0	Frascati	5 ± 2	33
^{197}Au	0.3 - 0.9	Lund	4 ± 2	25
	0.3 - 1.0	Frascati	$8. \pm 3$	33
^{238}U ^(a)	0.3 - 1.0	Frascati	10 ± 5	26

^(a) There is in the literature another measurement of the reaction $^{238}\text{U}(\gamma,n)^{237}\text{U}$ (Ref.28), but the reported values are 13 ± 5 mb in the energy range 0.2-0.5 GeV, and 4 ± 15 mb in the range 0.5-0.7 GeV.

unstable to proton emission. A small energy transfer to the rest nucleus during the photon interaction may easily excite ^{13}N which will then decay to the stable ^{12}C nucleus. No explanation has been found for the very high value of the cross section of the $^{103}\text{Rh}(\gamma, n)^{102}\text{Rh}$ reaction.

From the straight line fit, the following mass number dependence of the mean cross section $\bar{\sigma}_k$ was found

$$\bar{\sigma}_k = 0.104 A_t^{0.81} \text{ mb} , \quad (3)$$

with a correlation coefficient $r = 0.80$.

The good correlation we obtained in Ref. 40, between the logarithm of $\bar{\sigma}_k$ and the logarithm of A_t , strongly suggested that a linear dependence between these two variables should have been valid. As a consequence, a more refined least squares analysis was carried out by using the method of successive iterations with data rejection, i.e., after each iteration all those experimental points whose deviations from the calculated values were larger than twice the standard deviation were rejected. A total of four iterations were needed (for the 5th iteration, all the remaining points fell within twice the standard deviation), and only 20% of the experimental points having been rejected. A correlation coefficient $r = 0.94$ has been obtained in such a way. The deduced A_t -dependence of the calculated mean cross section per photon was found to be

$$\bar{\sigma}_F = (0.083 \pm 0.008) A_t^{(0.880 \pm 0.012)} \text{ mb}, \quad (4)$$

where σ_F denotes, henceforth, the value calculated from this fit.

Both Eqs.(3) and (4) give an A_t -dependence of $\bar{\sigma}(\gamma, n)$ of the type

$$\bar{\sigma} = a A_t^b , \quad (5)$$

with a b exponent greater than 2/3 (which would arise from a surface pion production model^{41,42}). A recent paper⁴³ correlates the trend of direct reactions with a volume production model^{44,45} and with more recent experimental data on pion photoproduction^{46,47}. The same conclu-

sions have been reached by Heimlich et al.⁴⁸ for the electroproduction of negative pions.

Figure 1 shows the trend of $\bar{\sigma}_F(\gamma, n)$ versus the target mass number, A_t , as calculated from Eq. (4). The straight line has been back-extrapolated up to $A_t = 1$ for reasons which will be clear in a next Section of this paper. The shaded area represents twice the error in the calculated $\bar{\sigma}_F$ values, as deduced from the fit error in the coefficients a and b . The Figure also reports the experimentally determined cross sections for the sake of comparison.

The relationship between $\bar{\sigma}_F$ and A_t , found by a simple statistical treatment of the available experimental data, will be soon compared with that obtained by a stochastic treatment of the interactions that lead to a (γ, n) reaction.

The next Section will describe the essentials of the Monte Carlo calculation which has been performed in order to obtain the cross section of the (γ, n) reactions.

3. USE OF THE MONTE CARLO METHOD IN THE STUDY OF THE (γ, n) REACTION

There exist in the literature several papers concerning the use of the Monte Carlo method for intranuclear cascades, initiated by high-energy incident protons on complex nuclei⁴⁹⁻⁵². In the absence of self-consistent theories covering the field of high-energy nuclear reactions, the Monte Carlo method turned out to be a powerful tool in obtaining a number of important nuclear parameters which characterize the post-cascade nucleus. However, in the case of photons as incident particles, only a few applications of this method have been developed, cf. Refs. 38, 39, 53, 54. As the elementary γ -nucleon (γ -quasideuteron) and the subsequent particle-nucleon collisions are at present well known, the Monte Carlo method is entirely adequate in estimating the yields of specific photo-reactions. In fact, Barashenkov et al.³⁹ were successful in carrying out a Monte Carlo calculation of photon-induced intranuclear cascades. A satisfactory agreement was observed between calculated and experimental data.




Fig.1 - The experimental (γ, n) cross sections versus the mass number of the target nucleus, A_t . The straight line is a least squares fit of the experimental points. The shaded area represents twice the statistical error. The dashed line represents an extrapolation up to $A_t = 1$. Experimental data are taken from: \triangle , Ref. 12; \blacktriangle , Ref.13; \blacktriangledown , Ref. 14 ; \square , Refs. 15,19,24,26; \triangledown , Refs. 17,18,20-23,25,27; \circ , Refs. 29, 30,32; \diamond , Ref. 31; \bullet Ref. 33.

In this Section, we describe briefly the successive steps of the Monte Carlo calculation used in order to obtain the probabilities of (γ, n) reactions for complex nuclei ranging between ^{12}C and ^{238}U , and for incident photons of intermediate energies (0.2-1.0 GeV).

3.1. The Nuclear Model

We assumed the target nucleus as described by a degenerate Fermi gas of nucleons confined within a spherically symmetric nuclear potential of radius $r_0 A_t^{1/3}$. Different values of r_0 were assumed according to the mass range of the target nucleus⁵⁵: $r_0 = 1.4$ fm for $A_t < 30$; $r_0 = 1.3$ fm for $30 < A_t < 100$; and $r_0 = 1.2$ fm for $A_t > 100$.

According to Serber's model³, a high-energy particle reaction can be described by two independent steps mechanism. In the first one, a rapid intranuclear cascade is initiated by the incoming particle, and successive interactions take place between individualized nucleons. In the course of the calculation, we assumed classical trajectories in describing the scattering between particles⁵⁶. During the fast cascade stage, there is a possibility for some particles (nucleon and/or mesons), as well as some nucleon aggregates (nuclear clusters)⁵⁷, to escape from the nucleus. Absorption of nucleons and reabsorption of mesons can also occur, thus resulting in a transfer of a fraction of the incident energy to the target nucleus. In the second step, the excited residual nucleus loses its energy by the evaporation of a number of particles until a cold spallation residual nucleus is reached⁵⁸. For heavy nuclei, fission can also take place as a mechanism of de-excitation competitive with particle evaporation.

In the present calculation, we have taken into consideration only the development of the fast cascade step, since it is expected that a (γ, n) reaction will take place during those intranuclear cascades which lead to residual nuclei with an excitation energy not so high as to permit the evaporation of other particles.

We considered the following primary interactions:

quasi-deuteron interaction $\gamma + (np) \rightarrow n + p$, (6)

$$\gamma + n \rightarrow n + \pi^0 \quad (7)$$

$$\gamma + n \rightarrow p + \pi^- \quad (8)$$

$$\gamma + p \rightarrow p + \pi^0 \quad (9)$$

$$\gamma + p \rightarrow n + \pi^+ \quad (10)$$

photomesonic interactions $\gamma + n \rightarrow n + \pi^+ + \pi^-$ (11)

$$\gamma + n \rightarrow n + \pi^0 + \pi^0 \quad (12)$$

$$\gamma + n \rightarrow p + \pi^- + \pi^0 \quad (13)$$

$$\gamma + p \rightarrow p + \pi^+ + \pi^- \quad (14)$$

$$\gamma + p \rightarrow p + \pi^0 + \pi^0 \quad (15)$$

$$\gamma + p \rightarrow n + \pi^+ + \pi^0 . \quad (16)$$

In writing the reactions (7-16), we have disregarded the intermediate steps via the formation of nucleon isobars, only taking into consideration the products of the decay of those isobars.

Being the mean-free-path of intermediate-energy photons in nuclear matter (~ 140 fm) large with respect to the nuclear sizes, we were led to consider the nucleus as transparent to photons and, therefore, look at each point of the whole nucleus as an equally probable primary interaction point source. For the sake of simplicity, a reduction of the spherical nucleus to a two-dimensional geometry was made⁴⁹.

As has been pointed out by Gabriel and Alsmiller³⁸, it is very difficult to evaluate the number of quasi-deuterons within the nucleus when this latter is subdivided into three distinct regions having different nucleon densities. On the other hand, such a subdivision would lengthen to a very great extent the computing time for the calculation of the paths travelled by the meson and recoil nucleon or by the photodisintegrated nucleon pair within the nucleus in their escaping.

From the above considerations, we decided to treat the nucleus within the framework of a constant nucleon density.

The photoproduced particles which result from the primary interaction can

yield different kinds of secondary reactions. We have taken into account the following ones



as the more representative in describing the cascade step.

3.2. Input Information

The total cross sections for the primary interactions were taken from Refs.1,59-62. In contrast with the assumption of Gabriel and Alsmiller³⁸ who considered the Levinger's factor, L , as being constant for all the nuclei under investigation, we assumed an A_t -dependence of L as indicated in Table 2. The angular distribution for the reactions (6) and (19) was considered isotropic in the CM-system. The differential cross sections for reactions (7-10) are those reported in Refs.59,63, while for reactions (17-20) they are those of Metropolis et al.⁵⁰. For interactions like (18), we have made the assumption that the total momentum was equally distributed between the produced particles in the CM-system. The mean-free-path of the various particles in the course of a cascade was calculated following Refs. 34,35. The total cross sections used in calculating the various nucleon-nucleon collision probabilities were those of Bertini⁵¹.

Table 2 - Dependence of Levinger's Factor, L , on the Mass Number, as Used in the Present Monte Carlo Calculation

Nucleus	¹² C	¹⁹ F	²³ Na	⁵⁵ Mn	¹⁰³ Rh	¹⁹⁷ Au	²³⁸ U
L	5.5	7	7	9	11	12	12

3.3. The Course of the Calculation

The calculation itself was performed following the general way described by Metropolis *et al.*⁵⁰. Only events of the type

$$\gamma + \frac{A}{Z}X \rightarrow \frac{A-1}{Z}X + n \quad , \quad (21)$$

leading to residual nuclei with an excitation energy below the appropriate cutoff energy (see Table 3) were recorded. As a result, we obtained the different probabilities for the above reactions as defined by the ratio between the number of iterations yielding one-neutron deficient residual nuclei and the total number of iterations. The standard treatment of each case, viz., a fixed incident photon energy on a fixed target nucleus, involved about ten thousand incident photons. About a hundred cases were analysed in the present calculation. The following target nuclei were selected: ¹²C, ¹⁹F, ²³Na, ⁵⁵Mn, ¹⁰³Rh, ¹⁹⁷Au and ²³⁸U. They

Table 3 - Values of the Cutoff Energy and Related Nuclear Characteristics Used in the Present Monte Carlo Calculation. (All Energies in MeV)

Nucleus	Fermi Energy		Average Binding Energy of Loosest Nucleon	Coulomb Energy at Surface	Cutoff Energy (*)
	Protons	Neutrons			
¹² C	24.6	24.6	7.7	2.3	34.5
¹⁹ F	23.7	25.4	7.9	3.1	35.6
²³ Na	23.9	25.3	8.3	3.6	36.5
⁵⁵ Mn	23.1	26.1	8.7	6.5	39.8
¹⁰³ Rh	26.1	30.9	8.6	10.4	47.5
¹⁹⁷ Au	28.9	37.7	7.9	16.1	57.3
²³⁸ U	28.2	38.4	7.5	17.6	58.4

(*) The cutoff energy was calculated as the average Fermi energy of neutrons and protons plus the average binding energy of the loosest nucleon plus the Coulomb energy at surface.

were chosen in order to give a moderately good coverage over complex nuclei of the periodic table.

The choice of the first interaction point, the type of interaction and, in the case of photomesonic interaction, the target nucleon, was performed following the general procedure described by Rudstam⁴⁹. The dynamics of successive collisions was treated relativistically in the usual manner^{64,65}. The distance of travel, d , of the various particles was determined by the relation

$$d = -\lambda(E_i) \log \xi, \quad (22)$$

where ξ is a random number in the interval 0-1 and $\lambda(E_i)$ is the mean-free-path of particle i with kinetic energy E_i in nuclear matter.

The calculation was carried out by using the IBM/370 computer of the CBPF. The general course of the calculation, described in details in the fluxogram of Fig.2, has been programmed for the cascade step of any type of photonuclear reactions. For simple (γ, n) reactions, two remarkable simplifications have been made. At incident energies higher than or equal to 0.2 GeV, the kinematics of the quasideuteron photodisintegration leads to a neutron and a proton both with kinetic energies larger than the cutoff energy of the struck nucleus. In view of this, they either escape the nucleus simultaneously or, if one or both are absorbed, give to the nucleus an energy sufficient to allow for the evaporation of other particles. In any case, an event different from (21) would be registered.

In the case of double-pion photoproduction, a large amount of energy is generally transferred to the nucleus and this will result in the emission of a number of particles during the evaporation stage. As a consequence, the contribution of this process to the (γ, n) probability can be assumed as almost negligible.




Fig.2 - Block diagram of the course of the Monte Carlo calculation of photonuclear cascades.

4. RESULTS OF THE MONTE CARLO CALCULATION

4.1. (γ, n) Reaction Probability

We define the (γ, n) reaction probability, $\Phi_n(k, A_t)$, as the ratio of the (γ, n) reaction cross section to the total inelastic cross section. Of course, this probability is a function of the photon energy k and the target mass number A_t .

The values of Φ_n have been obtained by means of the procedure already discussed in the Section 3.3 and are listed in Table 4, which also reports the mean values $\bar{\Phi}_n$, averaged over the energy range 0.3-1.0 GeV.

In Fig.3, the trends of Φ_n versus the photon energy for the nuclei ^{12}C and ^{238}U are shown. The curves exhibit very broad bumps centered around 0.5 GeV. The same behaviour is also observed for the other nuclei we have considered.

Plotted in Fig.4 are the mean values $\bar{\Phi}_n$, in the energy range 0.3-1.0 GeV, as a function of the target mass number A_t . The straight line through the calculated points is a least squares fit which gives an A_t -dependence of the form

$$\bar{\Phi}_n(k, A_t) = 0.234 A_t^{-0.14}, \quad (23)$$

with an estimated error in $\bar{\Phi}_n$ ranging between 3% (^{12}C) and 6% (^{238}U). By back-extrapolating the straight line, one obtains the value $A_t = 1$.

4.2. The (γ, n) Cross Section

Above the threshold of pion photoproduction, the (γ, n) cross section is related to the total inelastic cross section in the following way

$$\sigma(k, A_t) = \left[A_t \sigma_N \delta + L(N_t Z_t / A_t) \sigma_d \delta' \right] \Phi_n(k, A_t), \quad (24)$$

where the expression in brackets represents the total inelastic cross section, which accounts for the photomeson (the first term in the sum) and

4 - The (γ, n) Reaction Probabilities, $\Phi_n(k, A_t)$, as Obtained by the Present Monte Carlo Calculations (*)

(GeV)	^{12}C	^{19}F	^{23}Na	^{55}Mn	^{103}Rh	^{197}Au	^{238}U
0.20	0.007 ± 0.002	0.012 ± 0.003	0.012 ± 0.003	0.019 ± 0.004	0.025 ± 0.005	0.034 ± 0.005	0.038 ± 0.006
0.25	0.092 ± 0.009	0.079 ± 0.008	0.077 ± 0.008	0.062 ± 0.007	0.027 ± 0.005	0.052 ± 0.007	0.053 ± 0.007
0.30	0.152 ± 0.011	0.134 ± 0.010	0.130 ± 0.010	0.110 ± 0.009	0.096 ± 0.009	0.085 ± 0.008	0.085 ± 0.008
0.32	0.169 ± 0.011	0.147 ± 0.010	0.147 ± 0.010	0.116 ± 0.010	0.108 ± 0.009	0.091 ± 0.009	0.087 ± 0.008
0.35	0.162 ± 0.011	0.147 ± 0.010	0.149 ± 0.011	0.133 ± 0.010	0.116 ± 0.010	0.104 ± 0.009	0.101 ± 0.009
0.40	0.161 ± 0.011	0.141 ± 0.010	0.143 ± 0.010	0.154 ± 0.010	0.119 ± 0.010	0.106 ± 0.009	0.108 ± 0.009
0.45	0.180 ± 0.012	0.170 ± 0.011	0.170 ± 0.011	0.153 ± 0.011	0.135 ± 0.010	0.137 ± 0.010	0.139 ± 0.011
0.50	0.163 ± 0.014	0.147 ± 0.011	0.129 ± 0.011	0.136 ± 0.011	0.123 ± 0.010	0.114 ± 0.010	0.113 ± 0.011
0.55	0.173 ± 0.014	0.165 ± 0.012	0.166 ± 0.013	0.152 ± 0.012	0.145 ± 0.012	0.133 ± 0.012	0.129 ± 0.011
0.60	0.167 ± 0.014	0.163 ± 0.012	0.158 ± 0.013	0.139 ± 0.012	0.134 ± 0.012	0.119 ± 0.011	0.116 ± 0.011
0.65	0.158 ± 0.014	0.153 ± 0.013	0.149 ± 0.013	0.128 ± 0.012	0.123 ± 0.012	0.107 ± 0.011	0.107 ± 0.011
0.70	0.171 ± 0.014	0.163 ± 0.013	0.159 ± 0.013	0.139 ± 0.012	0.133 ± 0.012	0.114 ± 0.011	0.110 ± 0.011
0.80	0.167 ± 0.014	0.166 ± 0.013	0.164 ± 0.013	0.140 ± 0.012	0.125 ± 0.012	0.111 ± 0.011	0.107 ± 0.011
0.90	0.144 ± 0.014	0.137 ± 0.013	0.131 ± 0.013	0.114 ± 0.012	0.104 ± 0.012	0.085 ± 0.011	0.084 ± 0.011
1.00	0.148 ± 0.014	0.135 ± 0.012	0.135 ± 0.013	0.114 ± 0.012	0.108 ± 0.010	0.085 ± 0.010	0.082 ± 0.010
$\bar{\Phi}_n$ (**)	0.161 ± 0.013	0.151 ± 0.012	0.148 ± 0.012	0.132 ± 0.011	0.121 ± 0.011	0.106 ± 0.010	0.104 ± 0.010

(*) The quoted errors are statistical only.

(**) The symbol $\bar{\Phi}_n$ represents the mean value of $\Phi_n(k, A_t)$ in the energy range 0.3 - 1.0 GeV.




Fig.3 - The (γ, n) probability, $\Phi_n(k, A_t)$, versus incident photon energy. The points represent the results of the present Monte Carlos calculation. For the sake of simplicity, only the (γ, n) probabilities for ^{12}C and ^{238}U are shown. The curves are eye fits through the calculated points.




Fig.4 - Plot of the mean value of the (γ, n) probability, $\bar{\Phi}_n(k, A_t)$, in the energy range 0.3-1.0 GeV, versus the mass number of the target nucleus, A_t . The points have been calculated from the values listed in Table 4. The full line is a least squares fit of the calculated points. The dashed line represents an extrapolation up to $A_t = 1$. The symbol ✕ denotes the ratio of the cross section of the $\gamma + p \rightarrow n + \pi^+$ reaction to the total inelastic cross section, and ▶ stands for the ratio of the cross section of the $\gamma + n \rightarrow n + \pi^0$ reaction to the total inelastic cross section.

the quasideuteron interaction mechanisms. The quantities δ and δ' , which also depend on A_t and k , are factors related to the nuclear excitation, according to the type of interaction, and are very close to unity at energies below 1 GeV. The values of $a_N(k)$ have been taken from Ref. 66, and those of $\sigma_d(k)$ from Ref. 1.

By using the Φ_n probabilities of Table 4, we obtained the values of $\sigma(k, A_t)$ for (γ, n) reactions, hereafter denoted by σ_{MC} , listed in Table 5. In Fig.5, the trends of σ_{MC} are reported for ^{12}C and ^{238}U .

By least squares fitting the mean values $\bar{\sigma}_{MC}$ (0.3-1.0 GeV) taken from Table 5, the straight line shown in Fig.6 has been obtained. The A_t -dependence of $\bar{\sigma}_{MC}$ has the following expression:

$$\bar{\sigma}_{MC} = (0.063 \pm 0.006) A_t^{0.860 \pm 0.012} \text{ mb} . \quad (25)$$

Also in this case an exponent higher than 2/3 and very close to the exponent of Eq. (4) has been found.

5. DISCUSSION

5.1. Comments on the Treatment of Experimental Data

The relatively large number of experimental data on the (γ, n) reaction, covering the regions of light ($12 \leq A_t \leq 31$), medium-weight ($31 \leq A_t \leq 103$) and heavy ($103 < A_t \leq 238$) nuclei, allowed for a good statistical treatment which led to Eq. (4).

Only mean cross sections per photon were used, due to the lack of a large number of experimental values of cross sections per equivalent quantum at the different *Bremsstrahlung* energies for each target nucleus. Owing to this fact, a square shape of the *Bremsstrahlung* spectra has been assumed in calculating the mean cross sections per photons (see Eq.(2)) and consequently these cross sections are slightly overestimated (see also Section 2).

Table 5 - Calculated (γ, n) Reaction Cross Sections (mb)^(*)

$k(\text{GeV})$	^{12}C	^{19}F	^{23}Na	^{55}Mn	^{103}Rh	^{197}Au	^{238}U
0.20	0.12 ± 0.03	0.15 ± 0.05	0.17 ± 0.06	0.29 ± 0.10	0.75 ± 0.26	2.08 ± 0.73	0.81 ± 0.47
0.25	0.43 ± 0.12	0.62 ± 0.19	0.73 ± 0.22	1.55 ± 0.46	1.33 ± 0.40	5.00 ± 1.50	6.13 ± 1.70
0.30	0.94 ± 0.25	1.34 ± 0.33	1.61 ± 0.40	3.46 ± 0.86	5.77 ± 1.44	10.09 ± 2.52	12.14 ± 3.00
0.32	1.04 ± 0.25	1.49 ± 0.37	1.79 ± 0.45	3.57 ± 0.94	6.34 ± 1.58	10.50 ± 2.62	12.15 ± 3.00
0.35	0.89 ± 0.25	1.32 ± 0.33	1.62 ± 0.40	3.61 ± 0.90	5.99 ± 1.49	10.51 ± 2.62	12.30 ± 3.00
0.40	0.62 ± 0.17	0.88 ± 0.21	1.06 ± 0.25	2.90 ± 0.70	4.25 ± 1.02	7.42 ± 1.78	9.09 ± 2.00
0.45	0.51 ± 0.17	0.76 ± 0.18	0.93 ± 0.22	2.05 ± 0.49	3.44 ± 0.83	6.82 ± 1.64	8.32 ± 2.10
0.50	0.39 ± 0.09	0.56 ± 0.12	0.60 ± 0.13	1.56 ± 0.34	2.67 ± 0.59	4.82 ± 1.06	5.75 ± 1.24
0.55	0.44 ± 0.13	0.67 ± 0.15	0.82 ± 0.18	1.82 ± 0.40	3.28 ± 0.72	5.84 ± 1.28	5.80 ± 1.20
0.60	0.47 ± 0.13	0.70 ± 0.15	0.86 ± 0.19	1.83 ± 0.40	3.33 ± 0.73	5.66 ± 1.25	6.26 ± 1.20
0.65	0.45 ± 0.13	0.70 ± 0.15	0.82 ± 0.18	1.69 ± 0.37	3.04 ± 0.67	5.08 ± 1.12	6.10 ± 1.20
0.70	0.51 ± 0.14	0.77 ± 0.19	0.92 ± 0.23	1.91 ± 0.48	3.43 ± 0.86	5.63 ± 1.40	6.57 ± 1.20
0.80	0.50 ± 0.14	0.78 ± 0.19	0.94 ± 0.23	1.91 ± 0.48	3.20 ± 0.80	5.42 ± 1.11	6.31 ± 1.20
1.00	0.39 ± 0.11	0.56 ± 0.17	0.69 ± 0.20	1.38 ± 0.41	2.44 ± 0.73	4.88 ± 1.46	4.32 ± 1.30
$\bar{\sigma}_{\text{MC}}^{(**)}$	0.54 ± 0.13	0.80 ± 0.16	0.96 ± 0.20	2.00 ± 0.43	3.57 ± 0.75	6.28 ± 1.20	6.94 ± 1.20

(*) The cross sections listed in this Table were calculated from Eq.24. The quoted errors result from a combination of errors in the elementary cross sections and in the (γ, n) reaction probabilities (see Table 4).

(**) The symbol $\bar{\sigma}_{\text{MC}}$ represents the mean value of the cross sections in the energy range 0.3-1.0 GeV.




Fig.5 - Trends of the (γ, n) reaction cross sections as a function of incident photon energy. The points are taken from Table 5 (for the sake of clearness only the cross sections of ^{238}U and ^{12}C are shown). The full lines are eye fits through the calculated points. For comparison, it is also shown the experimental trend for ^{12}C as reported by Anderson *et al.*²⁷ (dashed line).





Fig.6 - A_t -dependence of the (γ, n) mean cross section, $\bar{\sigma}_{MC}$. The points are the calculated cross sections listed in Table 5. The straight line marked $A_t^{0.86}$ is a least squares fit of the calculated points. For the sake of comparison, the trends are also reported for a surface production model ($A_t^{2/3}$) and a volume production model (A_t^1).

In the case of the $^{12}\text{C}(\gamma, n) \text{ }^{11}\text{C}$ reaction, for which a larger number of experimental data were available, it was possible to determine the trend of σ_R , as a function of the photon energy, by solving Eq. (1) (see Ref. 27). However, the resolution of this integral equation gives strongly oscillating solutions due to its mathematical structure^{6,7}.


We also wish to point out that, by extrapolating Eq. (4) to $A_t = 1$, a cross section value of 0.083 mb is obtained, which is very close to the mean cross section values of the elementary processes $\gamma + p \rightarrow n + \pi^+$ ($\bar{\sigma} = 0.070$ mb) and $\gamma + n \rightarrow n + \pi^0$ ($\bar{\sigma} = 0.096$ mb). Such an agreement can be considered reasonable if we recall the large errors involved in the primary cross section measurements and in the coefficient of Eq. (4).

5.2. A Deeper Insight into the (γ, n) Reaction Probability

Equation (23) of Section 4.1 gives the A_t -dependence of the mean value of the (γ, n) reaction probability $\bar{\Phi}_n$. The coefficient 0.234 of Eq. (23) must represent the mean (γ, n) probability for a target having $A_t = 1$. For $A_t = 1$, two types of target have to be taken into consideration, i.e., either a proton target or an (idealized) neutron target, the corresponding interactions being therefore



and



In both cases, in fact, the photoproduction of a single neutron is obtained.

The mean probabilities of these processes are obtained from the mean ratios between the cross sections of each process and the total inelastic cross section per nucleon, via the photomesonic mechanism of interaction. These ratios are found to be 0.36 and 0.24, respectively, and are in good agreement within the limits of errors, which are quite large, with the value 0.234 arising from Eq. (23) (see Fig. 4). One should consider, though, that the first elementary process shown above (Eq. (26)) is effective in giving a direct (γ, n) reaction for $A_t = 1$ solely, since for $A_t > 1$ it may

contribute to the (γ, n) reaction only through an indirect way. On the contrary, the other process (Eq.(27)) will contribute in any case to the (γ, n) reaction. As a matter of fact, we found an extrapolated value which is very close to the mean calculated probability for the second process. This result is easily explained if one remembers the higher escaping probabilities of uncharged particles.

As has already been said, the (γ, n) probability Φ_n is a function of the target mass number and photon energy. Let us write this function as

$$\Phi_n(k, A_t) = \phi(A_t) \eta(k) , \quad (28)$$

i.e., as the product of two functions, each depending on one variable only. Further, let us assume $\phi(A_t)$ be of the form,

$$\phi(A_t) = A_t^\alpha .$$

In this way, Eq. (28) becomes

$$\Phi_n(k, A_t) = A_t^\alpha \eta(k) . \quad (29)$$

By using for $\bar{\sigma}(\gamma, n)$ the expression of Eq.(24), we obtain

$$\bar{\sigma}(\gamma, n) = \left\langle \left[A_t \sigma_N + L \frac{N_t Z_t}{A_t} \sigma_d \right] \Phi_n(k, A_t) \right\rangle . \quad (30)$$

Since $N_t Z_t / A_t$ is roughly approximated by $0.25 A_t$, and Levinger's factor by $L = 2.68 A_t^{0.28}$ (see Ref.40), Eq. (30) becomes

$$\bar{\sigma}(\gamma, n) = A_t \left\langle \left[\sigma_N + 0.67 A_t^{0.28} \sigma_d \right] \Phi_n(k, A_t) \right\rangle \quad (31)$$

and substituting $\Phi_n(k, A_t)$ from Eq. (29) gives

$$\bar{\sigma}(\gamma, n) = A_t^{\alpha+1} \left\langle \left[\sigma_N + 0.67 A_t^{0.28} \sigma_d \right] \eta(k) \right\rangle . \quad (32)$$

At this point, we can disregard the second term in the expression within

brackets, since it represents only 5% of σ_N for ^{12}C , and 12% for ^{238}U , thus obtaining

$$\bar{\sigma}(\gamma, n) \approx A_t^{\alpha+1} \langle \sigma_N(k) \eta(k) \rangle, \quad (33)$$

$\langle \sigma_N(k) \eta(k) \rangle$ being now a function of the photon energy only. By comparing Eq.(33) with Eq.(25), it can be written with a satisfactory degree of approximation as

$$\begin{aligned} \langle \sigma_N(k) \eta(k) \rangle &= 0.063 \text{ mb}, \\ \alpha + 1 &= 0.86, \end{aligned} \quad (34)$$

whence, $a = -0.14$.

Finally, Eq.(29) may be rewritten as

$$\Phi_n(k, A_t) = A_t^{-0.14} \eta(k) \quad (35)$$

from which

$$\eta(k) = \Phi_n A_t^{0.14}. \quad (36)$$

Values of $\eta(k)$ have been calculated for different nuclei, as a function of the photon energy, by means of Eq.(36), and by using for Φ_n the values listed in Table 4. At each energy, $\eta(k)$ was found to be approximately constant for all target nuclei under investigation, thus supporting the idea of taking η as a function of the photon-energy only.

In Fig.7, the trend of η/k is shown as a function of the photon-energy. By least squares fitting the points in the figure, a straight line has been drawn from which the following equation is obtained:

$$\eta(k) = 1.25 \times k \exp[-k/0.55], \quad (37)$$

k being expressed in GeV. The function η reaches its maximum at 0.55 GeV. This means that the (γ, n) probability should also reach a maximum at the




Fig.7 - The energy dependence of the (γ, n) reaction probability, $\eta(k)$. (For details see text). The points are calculated values of η/k . The straight line was obtained by a least squares fit. The error bars are statistical uncertainties.

same energy. This behavior has already been found for Φ_n , as shown in Fig.3. An explanation of such a trend may lie in the fact that the mean-free-path of the photoproduced particles increases from 0.3 GeV up to about 0.5 GeV, and the increased mean-free-path enhances the probability of a (γ, n) reaction via a direct process. Above about 0.5 GeV the double-pion photoproduction, instead, lowers considerably this probability, since this process certainly contributes to the total inelastic cross section but it does not contribute markedly to the (γ, n) process in view of the high excitation energy given to the nucleus.

The mean value of η between 0.3 GeV and 1.0 GeV turns out to be 0.237 which is in good agreement with the value 0.234 of Eq. (23) for $A_t = 1$ (see also Eq. (36)).

5.3. Comparison Between Experimental and Calculated (γ, n) Cross Sections

The most general relationship between the (γ, n) reaction cross section and the total inelastic cross section is given by Eq. (24), which is valid for any target nucleus and for energies above the giant resonance region.

By means of the Monte Carlo method, we succeeded in calculating the (γ, n) probability $\Phi_n(k, A_t)$ and thus in deducing the cross sections of (γ, n) reactions. These values, listed in Table 5, are shown in graphical form in Fig. 5, where the trends for ^{12}C and ^{238}U only are reported for the sake of clarity. Figure 5 also reports the trend of $\sigma(\gamma, n)$ for ^{12}C taken from Ref. 27. A satisfactory agreement between the calculated and experimental curves has been obtained.

A more simplified expression for $\sigma(\gamma, n)$ can be established by disregarding the quasideuteron contribution to the total inelastic cross section and by using Eqs. (35) and (37), which give

$$\sigma_C(\gamma, n) = 1.25 \sigma_N k \exp[-k/0.55] A_t^{0.86} \text{ mb}, \quad (k \text{ in GeV}), \quad (38)$$

where σ_C denotes the cross section calculated in this way

Mean values of the cross sections per photon of the (γ, n) reaction have been calculated by means of Eq. (4) (least squares fit of the experimental data), Eq. (24) (Table 4), and Eq. (25) (least squares fit of the Monte Carlo calculated points). We also calculated the means cross sections per photon by means of Eq. (38). The results are summarized in Table 6. From the analysis of Table 6, it can be concluded that only a few experimental values are not well reproduced by the calculated cross sections, namely, ^{14}N (Ref. 23), ^{55}Mn (Ref. 17), ^{75}As (Ref. 22), ^{103}Rh (Ref. 19) and ^{127}I (Ref. 14). In the other cases, experimentally determined cross sections are reproduced within a factor less than or equal to two.

The fact that the accuracy in predicting (γ, n) cross sections varies within a factor two might seem to be rather a bad one, but it should be remembered that the range of values covered is quite large (a factor of more than 25) and the same is true of the range of nuclear masses (10-240). Moreover, the experimentally determined cross sections are quite often very uncertain and the precision with which the four different equations (Eqs. (4), (24), (25) and (38)) predict (γ, n) cross sections is probably better than the uncertainty factors would indicate.

5.4. Mass Number-Dependence of the Cross Sections

A number of papers attempted to correlate the yield of pion photoproduction with the mass number or the target nucleus. Early measurements of charged pion photoproduction from nuclei showed an $A_t^{2/3}$ -dependence of the cross sections upon the target mass number A_t and this dependence was related^{41, 42} to a surface production mechanism (surface production model). Further experiments have shown, on the contrary, that the phenomenon actually exhibits an $A_t^{3/4}$ -dependence and this has led to a volume production model⁴³⁻⁴⁵ which also accounts for a more or less strong reabsorption of pions within the nucleus.

More recent experiments⁴⁶⁻⁴⁸ seem to confirm the predictions of the volume production model. Shramenko et al.⁴⁷, by using known values of nuclear transparencies^{34, 35}, found an 0.88-dependence for the photoproduction of π^+ - and π^- -mesons.

Table 6 - Comparison Between Experimental and Calculated Mean (γ, n) Reaction Cross Sections (mb) at Intermediate Energies

Target Nucleus	Experimental Cross Section		Calculated Cross Sections			
	Ref.	σ_k	$\bar{\sigma}_F$ (Eq.4)	$\bar{\sigma}_{MC}$ (Table 5, Eq.24)	$\bar{\sigma}_{MC}$ (Eq.25)	$\bar{\sigma}_c$ (Eq.38)
^{12}C	12	0.85 ± 0.10				
	13	0.4 ± 0.1				
	15	1.18 ± 0.20	0.74 ± 0.09	0.54 ± 0.13	0.53	0.52
	21,27	0.6 ± 0.1				
	32	0.9 ± 0.1				
^{14}N	23	0.11 ± 0.05	0.85		0.61	0.59
^{16}O	23	0.6 ± 0.1				
	29	1.27 ± 0.20	0.95		0.68	0.66
^{19}F	23	0.42 ± 0.10				
	24	1.33 ± 0.20	1.11 ± 0.15	0.80 ± 0.16	0.79	0.77
	30	1.30 ± 0.10				
^{23}Na	30	1.60 ± 0.20	1.3 ± 0.2	0.96 ± 0.20	0.93	0.90
^{31}P	23	0.9 ± 0.1				
	33	1.6 ± 0.1	1.70		1.21	1.17
^{52}Cr	33	2.5 ± 0.5	2.69		1.88	1.83
^{55}Mn	17	7.7 ± 0.5				
	31	4.2 ± 0.6	2.8 ± 0.4	2.0 ± 0.4	1.98	1.91
	33	3.5 ± 0.5				
^{59}Co	18	3.3 ± 0.5				
	33	3.0 ± 0.5	3.00		2.10	2.04
^{75}As	22	8.7 ± 1.0				
	33	4 ± 1	3.71		2.58	2.51
^{103}Rh	19	13 ± 5	4.90 ± 0.75	3.57 ± 0.75	3.39	3.35
^{127}I	14	10 ± 6				
	20	4 ± 2	5.89		4.06	3.94
	33	5 ± 2				
^{197}Au	25	4 ± 2	9 ± 1	6.28 ± 1.20	5.92	5.74
	33	8 ± 3				
^{238}U	26	10 ± 5	10 ± 2	6.94 ± 1.20	6.97	6.76

Above about 0.2 GeV, a (γ, n) reaction may occur via the photomesonic interaction mechanism only, the quasideuteron mechanism being energetically ineffective in producing a (γ, n) reaction.

Taking into account the primary interactions (7-16) listed in Section 3.1, it is clear that both neutrons and protons do contribute to a (γ, n) reaction and thus the cross section of this process must depend on the mass number A_t and not on the neutron number N_t only.

By means of quite different method, we obtained the following mass number dependences of $\bar{\sigma}(\gamma, n)$: $A_t^{0.88}$ from the statistical treatment of the experimental data, and $A_t^{0.86}$ from the Monte Carlo calculation. The conclusion can be therefore drawn that also in this case, the volume production model is adequate in describing the mechanism of interaction leading to a (γ, n) reaction. Such a conclusion is clearly shown by the trends shown in Fig. 6.

6. CONCLUSIONS

We wish to summarize briefly the most interesting results that have been obtained in the course of this work.

i) In spite of the fact that the Monte Carlo calculation has been carried out by using monoenergetic photons as incident particles, a good agreement has been found with the experimental mean cross sections deduced from *Bremsstrahlung* yields.

ii) The cross sections of (γ, n) reactions increase with increasing target mass number values, A_t , in the energy range 0.3-1.0 GeV.

iii) The (γ, n) reaction probability decreases with increasing target mass number values, A_t , in the same energy range.

iv) The (γ, n) reaction probability reaches a maximum at about 0.55 GeV.

v) The quasideuteron mechanism of interaction does not contribute to the (γ, n) reaction at energies above 0.2 GeV.

vi) Double- and multiple-pion photoproduction play an almost negligible role in producing a (γ, n) reaction.

vii) Both neutrons and protons are sources of primary interactions which may lead to a (γ, n) process.

The present paper is only a part of a systematic work aiming to study in detail the (γ, xn) ($x \geq 1$) reactions, as well as the $(\gamma, yp xn)$ (spallation) reactions, with the aid of the Monte Carlo calculation. Future work will be devoted to the study of the (γ, xn) ($x > 1$) reactions and the results will be the subject of future communications.

The authors wish to acknowledge the tireless collaboration of the computer staff of the *Centro Brasileiro de Pesquisas Físicas*. J. B. Martins and O.A.P. Tavares wish to express their appreciation of the very kind hospitality they received from the *Istituto di Chimica Generale ed Inorganica dell'Università*, Roma.

REFERENCES

1. J.S. Levinger, Phys. Rev. 84, 43 (1951); 97, 970 (1955).
2. I. Reff, Phys. Rev. 91, 150 (1953).
3. R. Serber, Phys. Rev. 72, 1114 (1947).
4. M.L. Goldberger, Phys. Rev. 74, 1269 (1948).
5. C.E. Roos and V.Z. Peterson, Phys. Rev. 124, 1610 (1961).
6. G.C. Baldwin and G.S. Klaiber, Phys. Rev. 73, 1156 (1948).
7. B.M. Spicer, Suppl. Nuovo Cimento 2, 243 (1964).
8. B. Bülow and B. Forkman, Lund Preprint, to be published by A. B. Atomenergi, Studsvik, Sweden (ed. D. Brune).
9. B.L. Berman and S.C. Fultz, Rev. Mod. Phys. 47 (1975).
10. K. Tesch, Nucl. Instr. Methods 95, 245 (1971).
11. W.C. Barber, W.D. George and D.D. Reagan, Phys. Rev. 98, 73 (1955).
12. A. Masaike, J. Phys. Soc. Japan 19, 427 (1964).
13. Yu. P. Antuf'ev, I.I. Miroshnichenko, V.I. Noga, P.V. Sorokin, Sov. J. Nucl. Phys. 6, 312 (1968).
14. G.G. Jonsson, B. Forkman, K. Lindgren, Phys. Lett. 26B, 508 (1968).
15. V. di Napoli, F. Dobici, O. Forina, F. Salvetti and H.G. de Carvalho, Nuovo Cimento 55B, 95 (1968).
16. H.G. de Carvalho, V. di Napoli, D. Margadonna, F. Salvetti and K. Tesch, Nucl. Phys. A126, 505 (1969).

17. G. Andersson, B. Forkman and B. Friberg, University of Lund Report N^o LUNP 6901, January 1969, p.11 (unpublished).
18. G. Andersson and B. Forkman, University of Lund Annual Report 1969, Sect. 5-A: Id.
19. V. di Napoli, D. Margadonna, F. Salvetti, H.G. de Carvalho and J. B. Martins, Lett. Nuovo Cimento 1, 308 (1969).
20. G.C. Jonsson, K. Lindgren and B. Forkman, University of Lund Report N^o LUNP 6901, January 1969, p. 14 (unpublished).
21. G. Hyltén, Nucl. Phys. A158, 225 (1970).
22. G. Andersson and B. Forkman, University of Lund Annual Report 1970, p.48.
23. B. Friberg, G. Andersson, and B. Forkman, Nucl. Phys. A171, 551 (1971).
24. V. di Napoli, D. Margadonna, F. Salvetti, H.G. de Carvalho and J. B. Martins, Nucl. Instr. Methods 93, 77 (1971).
25. K. Lindgren and G.G. Jonsson, Nucl. Phys. A166, 643 (1971).
26. V. di Napoli, F. Salvetti, H.G. de Carvalho and J. B. Martins, Lett. Nuovo Cimento 1, 538 (1971).
27. G. Andersson, I. Blomqvist, B. Forkman, G. G. Jonsson, A. Järund, I. Kroon, K. Lindgren, B. Schröder and K. Tesch, Nucl. Phys. A197, 44 (1972).
28. B. Schröder, B. Nordgren and A. Alm, University of Lund Report N^o LUNP 7206, March 1972, p. 17.
29. V. di Napoli and M. L. Terranova, Gazz. Chim. Ital. 203, 551 (1973).
30. F. Salvetti, G. Aurisicchio, V. di Napoli, M. L. Terranova, H. G. de Carvalho and J. B. Martins, Gazz. Chim. Ital. 103, 1003 (1973).
31. H. G. de Carvalho, J. B. Martins, O.A.P. Tavares, V. di Napoli and F. Salvetti, *Proc. Int. Conference on Photonuclear Reactions and Applications*, Pacific Grove (Cal.), March 26-30, 1973, Section 8C5-1.
32. V. di Napoli, M.L. Terranova, H.G. de Carvalho and J.B. Martins, Gazz. Chim. Ital. 104, 463 (1974).
33. V. di Napoli, F. Salvetti, M.L. Terranova, H.G. de Carvalho, J.B. Martins, and O.A.P. Tavares, Gazz. Chim. Ital. 105, 67 (1975).
34. H.G. de Carvalho, J.B. Martins, O.A.P. Tavares, R.A.M.S. Nazareth and V. di Napoli, Lett. Nuovo Cimento 2, 1139 (1971).
35. H.G. de Carvalho, J.B. Martins, O.A.P. Tavares, R.A.M.S. Nazareth and V. di Napoli, Lett. Nuovo Cimento 4, 365 (1972).
36. G. Andersson, B. Forkman and B. Friberg, University of Lund Report N^o LUNP 7010, September 1970.

37. G. Andersson, B. Forkman and B. Friberg, *Nucl. Phys.* **A171**, 529 (1971).
38. T.A. Gabriel and R.G. Alsmiller, Jr., *Phys. Rev.* **182**, 1035 (1969).
39. V.S. Barashenkov, F.G. Gereghi, A.S. Iljinov, G.G. Jonsson and V. D. Toneev, *Nucl. Phys.* **A231**, 462 (1974).
40. V. di Napoli, F. Salvetti, M.L. Terranova, H.G. de Carvalho, J.B. Martins and OAP. Tavares, *Gazz. Chim. Ital.* **105**, 317 (1975).
41. R.R. Wilson, *Phys. Rev.* **86**, 125 (1952).
42. S.T. Butler, *Phys. Rev.* **87**, 1117 (1952).
43. V. di Napoli, *Lett. Nuovo Cimento* **12**, 609 (1975).
44. J.R. Waters, *Phys. Rev.* **113**, 1133 (1959).
45. W.M. McClelland, *Phys. Rev.* **123**, 1423 (1961).
46. GN. Oudkin, V.N. Eponeshnikov, Yu. F. Krechetov and V.A. Tryasuchev, *Krakov Physico-Technical Institute Report N? KFTI 73-31, p.66 (1973) (in Russian)*.
47. B.I. Shramenko, I.A. Grishaev, V.I. Nikiforov and G.D. Pugachev, *Kharkov Physico-Technical Institute, Report N? KFTI 73-31, p.74 (1973) (in Russian)*.
48. F.H. Heimlich, G. Huber, E. Rössle, P. David, H. Mommsen and D. Wegener, *Nucl. Phys.* **A267**, 493 (1976).
49. G. Rudstam in "Spallation of Medium-Weight Elements", Doctoral Thesis, NP 6191, University of Uppsala, Uppsala (1956).
50. N. Metropolis, R. Bivins, M. Storm, A. Turkevich, J.M. Miller, and G. Friedlander, *Phys. Rev.* **110**, 185 (1958); N. Metropolis, R. Bivins, M. Storm, G.M. Miller, and G. Friedlander, *Phys. Rev.* **110**, 204 (1958).
51. H.W. Bertini, *Phys. Rev.* **188**, 1711 (1969).
52. V.S. Barashenkov, H.W. Bertini, K. Chen, G. Friedlander, G. D. Harps, A.S. Iljinov, J.M. Miller and V.D. Toneev, *Nucl. Phys.* **A187**, 531 (1972).
53. K.K. Gudima, A.S. Iljinov, and V.D. Toneev, *JINR communications* **P2-4661, P2-4808, Dubna (1969)**.
54. N.L. Emetz, G. Ya. Lubarsky, Yu. N. Ranyuk, and P.V. Sorokin, *Kharkov Physico-Technical Institute Report N? KFTI 72-37 (1972) (in Russian)*.
55. L.R.B. Elton, *Introductory Nuclear Theory*, Pitman, London (1965).
56. P.A. Benioff, *Phys. Rev.* **119**, 324 (1960).
57. V. di Napoli, F. Salvetti, M.L. Terranova, H.G. de Carvalho, and J.B. Martins, *Phys. Rev.* **C8**, 206 (1973).
58. V. Weisskopf, *Phys. Rev.* **52**, 295 (1937).

59. J.T. Beale, S.D. Ecklund and R.L. Walker, Report CTSL-42, CALT 68-108 (1968).
60. H.G. Hilpert *et al.*, ~~ABHM~~ Collaboration, Nucl.Phys. B8, 535 (1968).
61. C. Betourne, J.C. Bizot, J. Perez y Jorba, D. Treille, Phys. Lett. 24B, 590 (1967). See also other references quoted therein.
62. ~~ABHM~~ Collaboration, Phys. Rev. 175, 1669 (1968).
63. P. Spillantini and V. Valente, Report CERNHERA 70.1 (1970)
64. K.G. Dedrick, Rev. Mod. Phys. 34, 429 (1962).
65. J.D. Jackson in *Classical Electrodynamics*, John Wiley & Sons, London (1962).
66. M. Damashek and F.J. Gilman, Phys. Rev. D1, 1319 (1970).
67. D.L. Phillips, Journal ACM. 9 84 (1962).